

Publisher: Asian Economic and Social Society
ISSN (P): 2304-1455, ISSN (E): 2224-4433
Volume 3 No. 4 April 2013.

Asian Journal of Agriculture and
Rural Development



Effects of Variety and Fertilizers on Number of Grains/Cob of Maize in Asaba Area of Delta State

Enujeke E. C. (Department of Agronomy, Delta State University, Asaba Campus, Nigeria)

Citation: Enujeke E. C. (2013) “Effects of Variety and Fertilizers on Number of Grains/Cob of Maize in Asaba Area of Delta State”, Asian Journal of Agriculture and Rural Development, Vol. 3, No. 4, pp. 215-225.



Effects of Variety and Fertilizers on Number of Grains/Cob of Maize in Asaba Area of Delta State

Abstract

This study carried out in the Teaching and Research Farm of Delta state University, Asaba Campus from March 2008 to June, 2010 to evaluate the effects of variety, organic manure and inorganic fertilizer on number of grain/cob of maize. The experiment was carried out in a Randomized Complete Block Design (RCBD) replicated three times in a factorial layout. Four different rates of poultry manure, cattle dung and NPK 20:10:10 fertilizer were applied to three different maize varieties sown at 75cm x 15 cm and evaluated for number of grains/cob. The result obtained indicated that hybrid variety, 9022-13, which had 5090, was superior. Also, plants that received inorganic fertilizer were outstanding in number of grains/cob with values of 441.5 in 2008 and 506.0 in 2009 followed by the plants received poultry manure (444.0 in 2008 and 468.0 in 2009). Based on the rates of application of manure/inorganic fertilizer, plants that received 450kg ha^{-1} of NPK 20:10:10 fertilizer were outstanding in numbers of grains/cob of maize with values of 532.8 in 2008 and 570.8 in 2009. The results of interaction showed that variety, manure type and rates of application were significantly ($p < 0.05$) different. Based on the findings of this study, it is recommended that (i) Hybrid variety 9022-13 should be grown in Asaba area. (ii) NPK 20: 10:10 mineral fertilizers should be applied at the rate of 450kg ha^{-1} to enhance maize yield.

Author(s)

Enujeke E. C.

Department of Agronomy, Delta State University, Asaba Campus, Nigeria

Keywords: Organic manure, Inorganic fertilizer, Number of grains/cob, Asaba, Delta State

Introduction

Maize (*Zea mays* L) is one of the major cereal crops grown in the humid tropics and Sub-Saharan Africa. It is a versatile crop and ranks third following wheat and rice in world production as reported by Food and Agriculture Organization (FAO, 2002). Maize crop is a key source of food and livelihood for millions of people in many countries of the world. It is produced extensively in Nigeria, where it is consumed roasted, baked, fried, pounded or fermented (Agbato, 2003). In advanced countries, it is an important source of many industrial products such as corn sugar, corn oil, corn flour, starch, syrup, brewer's grit and alcohol (Dutt, 2005). Corn oil is used for salad, soap-making and lubrication. Maize is a major component of livestock feed and it is

palatable to poultry, cattle and pigs as it supplies them energy (Iken *et al.* 2001). The stalk, leaves, grain and immature ears are cherished by different species of livestock (Dutt, 2005).

In spite of the increasing relevance and high demand for maize in Nigeria, yield across the country continues to decrease with an average of about 1 t/ha which is the lowest African yield recorded (Fayenisin, 1993).

The steady decline in maize yield can be attributed to:

1. Rapid reduction in soil fertility caused by intensive use of land and reduction of fallow period as reported by Directorate of Information and

- Publications of Agriculture (DIPA, 2006).
2. Failure to identify and plant high yielding varieties most suited or adapted to each agro-ecological zone (Kim, 1997; Olakojo *et al.*, 1998).
 3. Use of inappropriate plant spacing which determines plant population and final yield (Zeidan *et al.*, 2006).
 4. Negligence for soil amendment materials such as organic manure and inorganic fertilizers which improve soil condition and enhance crop yield.

Sonetra (2002) suggested that subsistence farmers should apply organic manure directly to the soil as a natural means of recycling nutrients in order to improve soil fertility and yield of crops. Manures and fertilizers are the life wire of improved technology contributing about 50 to 60% increase in productivity of food grains in many parts of the world, irrespective of soil and agro-ecological zone (DIPA, 2006). Reijnties *et al.* (1992) and Adepetu (1997) remarked that the downward trend in food production should prompt farmers to amend the soil with different materials in order to enhance growth and yield of crops. Several organic materials such as cattle dung, poultry dropping, pig dung and refuse compost have been recommended to subsistence farmers in West Africa as soil amendments for increasing crop yield (Sobulo and Babalola, 1992; Ismail *et al.*, 1990; Olayinka, 1996 and Olayinka *et al.*, 1998).

Municipal wastes were reported to have reduced soil temperature, increased soil water, nutrient status and the yield of maize in temperate soils (Movahedi *et al.*, 2000). Cattle dung has been reported to contain 0.3 – 0.4 % N, 0.1 – 0.2 % P and 0.1 – 0.3 % K (Subedi and Gurung, 1991). According to Adekunle *et al.* (2005) cattle dung applied at the rate of 10 t/ha to cowpea resulted to increased plant height, leaf area, pod number, pod weight as well as improved soil structure in a mixed farming system. Stefan (2003) indicated that fresh poultry dropping contain 70% water, 1.4% N, 1.1% P₂O₅ and 0.5% K₂O while dried poultry manure contains 13% water, 3.6% N, 3.5% P₂O₅ and 1.6% K₂O. Ayodele (1993) reported that inorganic fertilizers are known to influence the quantity and yield of maize. Iken and Anusa (2004) recommended an

optimum plant population of 53,333 plants/ha for maximum yield of maize. Their report indicated that this is obtainable using a spacing of 75cm x 25cm at 1 plant per stand or 75cm x 50cm at 2 plants per stand. Azam *et al.* (2007) reported that spacing of 75cm x 35cm resulted in increased grain yield of maize while 75cm x 15cm gave maximum cob weight. Similar report by Alessi and Power (2004) revealed that maize cob weight decreased with increased plant population.

At present, there are no recommended standards taking into consideration the different combinations of such cultural practices as varietal selection, rates of appropriate organic manure and mineral fertilizer which interplay to influence yield and optimal performance of maize in Asaba area of Delta State. Against this background, the broad objective of this study, therefore, was to: identify variety of maize most suited or adapted to Asaba area and the appropriate fertilizer types and rates for the variety.

The specific objectives were to:

- (i) identify the best variety for Asaba area.
- (ii) determine the effects of NPK (20:10:10) mineral fertilizer, poultry manure, cow dung on number of grains/cobs maize.

Materials and Methods

Site Description

The study was carried out in the Teaching and Research Farm of Delta State University, Asaba Campus from March to December 2008 and repeated between March and December, 2009. Asaba is located at latitude 06°14'N and longitude 06°49'N of the equator. It lies in the tropical rainforest zone dominated by mangrove, fresh water, swamps, humid forests and secondary vegetation (NEST, 1991). Its climate is influenced by the movement of the Inter-Tropical Discontinuity (ITD). The IDT is made up of two wind systems namely the moisture-laden South-West monsoon from the Atlantic Ocean and the dry cold North-East trade wind from the Sahara desert. The South-West Trade wind most significantly determines the climate condition of Asaba area of Delta State. Asaba is characterized by raining season between April and October,

with annual mean-rainfall of 1500mm and 2000mm maximum. The distribution is bimodal with peak in July and September, coupled with a period of low precipitation in August. Mean temperature is 23.8°C with 37.3°C as maximum. Relative humidity is 77.2%, the mean monthly soil temperature at 100m depth is 20.3 °C, while sunshine stands at 4.8 bars (Meteorological Office, Asaba, 2003).

Pre-planting Soil Analysis

Representative surface soils (0-20cm) were sampled with a tubular sampling auger. These soil samples were air-dried at room temperature for 5 days and crushed to pass through a 2mm mesh sieve. Sub-samples from the bulked soil sample were further grounded to pieces to pass through 100mm-mesh sieve for the determination of organic matter. The rest samples were then analyzed to determine the physical and chemical properties of the soil. The analysis was done at Delta State University, Asaba campus.

Analytical Procedure

Physical Properties

Particle Size Distribution: Particle size distribution was analyzed using the Bouyoucos hydrometer method in which 0.5 N Sodium hexameta-phosphate was used as dispersant (Landor, 1991).

Bulk Density: The bulk density (Bd) was determined by Core-method.

Particle Density: This was determined by pycnometer or specific gravity bottle method as described by Bowles (1992).

Chemical Properties

Soil pH: This was determined in soil: water suspension (1:1) using glass electrode pH-meter as described by Mclean (1982).

Organic Carbon: This was determined using the wet oxidation method of Walkley and Black (Walkley and Black, 1945).

Total Nitrogen: This was determined using the modified K. Jeldah distillation method as described by Landor (1991).

Exchangeable Cations (EC) and Effective Cation Exchange Capacity (ECEC): Exchangeable cations were determined by extracting the cations with IN ammonium

acetate (IN, NHOAC) as displacing solution, buffered at pH₇ as described by Brady and Weils (1999). The extract was then determined electrochemically using atomic absorption spectrophotometry. The effective cation exchange capacity (ECEC) was calculated as the sum of exchangeable bases (Ca, Mg, K and Na) and exchangeable Al and H expressed in cmol/kg⁻¹ of soil.

Exchangeable Acidity: This was determined by titration method as described by Juo (1981). The exchangeable H⁺ and Al³⁺ were then expressed in cmol/kg⁻¹ of soil

Aim of the Experiment

The experiment was aimed at testing the response of three selected maize varieties to different rates of organic manure and inorganic fertilizer (NPK 20:10:10)

Experimental Design

The study was carried out in Randomized Complete Block Design (RCBD), replicated three times in a factorial layout. The factors were three sources of nutrients:- poultry manure (PM), cattle dung (CD), inorganic fertilizer (NPK 20:10:10). The different rates of PM were 0, 10, 20, 30t/ha, CD were 0, 10, 20, 30t/ha and NPK 20:10:10 were 0, 150, 300, 450kg/ha.

Agronomic Practices

Among the agronomic practices carried out were land preparation/plot layout, planting, application of treatments, weeding.

Land Preparation and Plot Layouts

The land was ploughed and harrowed using tractor. Three blocks (or replicates) consisting of 36 blocks each were layout, each block will measure 2.6m x 2.25m and was separated from one another with a space of 0.5m. Alley pathways of 1m separated one block from another, and the total number of plots laid out in the entire experiment was 108.

Planting

Maize seeds were sown on the plots at the rate of 1 seed per hole at a depth of 2-3cm, using 75cm x 15cm spacing as indicated by the first experiment.

Procurement and Application of Organic Manure and NPK 20:10:10 Fertilizer

Well-decomposed cattle dung was collected from cattle pen area, while poultry droppings were obtained from the battery cage system of poultry management of Delta State University, Asaba Campus. This organic manure was analyzed to determine their nutrient contents. NPK 20:10:10 fertilizer was obtained from Delta Agricultural Procurement Agency (DAPA), Ibusa. These amendment materials were incorporated into the plots according to the treatment as suggested by Olanikan (2006).

Weeding: Weeding was done three times using hoe.

Data Collection: Fourteen middle stands were used as sample population for data collection. Data collected was number of grains/cob of maize. This was done by direct counting after harvesting and shelling at the end of the sixteenth week from sowing.

Analysis of Soil Sample: Soil samples were collected from each experimental plot and analyzed for their content of N P and K.

Statistical Analysis: Data collected were subjected to analysis of variance (ANOVA) and means were separated with Duncan Multiple Range Test (DMRT) according to Wahua (1999).

Results

Soil Physico-chemical Properties of the Experimental Site:

The pre-physico-chemical properties of the experimental site are shown in Table 1. The result showed predominantly sand at the surface and this tends to decrease with depth of profile. Texturally, the soil of the experimental site is classified as sandy loam. The soil is acidic with pH of 6.2 in H₂O and 5.6 in CaCl. The organic matter content and total nitrogen were low with values of 1.22 gkg⁻¹ and 0.113 gkg⁻¹. The available P was high with value of 26.5 mgkg⁻¹. The exchangeable cations (Ca, Mg, Na and K) were equally low in status with values of 2.6cmolkg⁻¹ for Ca²⁺ and 0.9 cmolkg⁻¹ for Mg²⁺. The value obtained for Na⁺ was 0.57 cmolkg⁻¹, which was moderate while that for K⁺ was 0.08cmolkg⁻¹, which was low. The CEC was 4.15, while ECEC was 5.6cmolkg⁻¹, which were generally low. The exchangeable acidity was only trace for Al³⁺ and characteristically low for H⁺ with a value of 1.4 cmolkg⁻¹.

Nutrient Content (%) of Organic Manure Used in the study

The nutrient content of organic manure (poultry manure and cattle dung) used in the study is shown in Table 31. The values of N, P and K in poultry manure were significantly (P<0.05) higher than their values in cattle dung. With respect to N, poultry manure had 1.6% against cattle dung which was 0.4%. Also, poultry manure had 0.6% P while cattle dung had 0.2% P. The values for K were 0.8% in poultry manure, while it was 0.3% in cattle dung.

Table 1: Physico-chemical Properties of Experimental Site

Soil Property		Value	Interpretation
Particle Size Distribution (%)			
Coarse sand		38	
Fine sand		41	
Silt		9	
Clay		12	
Texture			Sandy loam
Ph	H ₂ O	6.2	Acidic
	CaCl	5.6	Acidic
Organic	Carbon gkg ⁻¹	0.71	
Organic	Matter gkg ⁻¹	1.22	Very low
Total	Nitrogen gkg ⁻¹	0.113	Low
Available	P (ppm)	26.5	
Exchangeable bases (cmol/kg ⁻¹)			
	Na ⁺	0.57	Moderate

	K ⁺	0.08	Very low
	Ca ²⁺	2.60	Low
	Mg ²⁺	0.90	Low
Cation Exchange Capacity		4.15	
Exchangeable acidity (cmol/kg ⁻¹)			
	Al ³⁺	Trace	
	H ⁺	1.4	
Effective cation Exchangeable capacity (Cmol/kg ⁻¹)		5.6	

Table 2: Nutrient (%) of Organic Manure used ion the Study

Parameter	Nutrient Content (%)		
	N	P	K
PM	1.6 _a	0.6 _a	0.8 _a
CD	0.4 _b	0.2 _b	0.3 _b

Legend: PM = Poultry Manure, CD = Cattle Dung, N = Nitrogen, P = Phosphorus

Effect of Variety, Organic Manure and Inorganic Fertilizer on Number of Grains/Cob of Maize

The response of number of grains/cob of maize to variety, organic manure and inorganic fertilizer is shown in Table 3. There were significant differences in the number of grains/cob of maize. With respect to varietal performance in 2008, hybrid variety 9022-13 had the highest number of grains/cob (526.6), followed by open-pollinated variety BR9922-DMRSRF₂ (474.0) Agbor local variety had the lowest number of grains/cob (340.0). The order of highest number of grains/cob with respect to the varieties sown was 9022-13 > BR9922-DMRSRF₂ > Agbor local. This was also the trend in 2009. Plants that received inorganic fertilizer NPK 20:10:10 had the highest number of grains/cob (506.0) followed by plants that received poultry manure (468.0). Plants that received cattle dung had the lowest

number of grains/cob. The order of superiority in the number of grains/cob based on source of nutrient was inorganic fertilizer NPK 20:10:10 > Poultry manure > cattle dung. Plants that received higher rates of fertilizer or manure had higher number of grains/cob than plants that received lower rates. The order of highest number of grains/cob based on rate of manure in tons/hectare was 30 > 20 > 10 > 0 while that of fertilizer in kg/ha was 450 > 300 > 150 > 0. Similar results were also obtained in 2009. The result of interaction (Table 4) showed that variety, manure type, rate, variety * rate were significantly (P<0.05) different and affected changes in number of grains/cob of maize in 2008 and 2009, while type * rate were not significantly (P>0.05) different and could not have affected changes in number of grains/cob of maize in both years of evaluation.

Table 3: Effects of Variety, Organic Manure and Inorganic Fertilizer on Number of Grains per Cob of Maize in 2008 and 2009

	No. of grains/cob		
	2008	2009	Mean
Variety			
9022 – 13	509.0 _a	526.6 _a	517.8 _a
BR9922-DMRSF ₂	410.0 _b	474.0 _b	242.0 _b
Agbor Local	343.3 _c	430.0 _c	386.6 _c
Nutrient Source			
PM	440.0 _a	468.0 _b	454.0 _a
CD	380.7 _b	458.0 _b	419.3 _b
IF	441.5 _a	506.0 _a	473.7 _a
Rates of application (tons or kg/ha)			
0	332.6 _g	394.0 _d	363.3 _e
manure (t/ha^{-1})	10	382.6 _f	423.0 _c
	20	434.6 _d	482.0 _b
	30	494.8 _b	542.0 _a
IF (kg/ha^{-1})	150	409.1 _e	496.6 _b
	300	473.3 _c	558.0 _a
	450	532.8 _a	570.8 _a

Means with the same letter(s) under the same column are not significantly different ($P \leq 0.05$) using Duncan Multiple Range test (DMRT).

Table 4: Effects of Variety, Manure and Inorganic Fertilizer on Number of Grains per Cob of Maize in 2008 and 2009

Variety	Nutrient Source	Rate	No. of Grain/cob	No. of Grain/cob
9022-13	PM	0	408.0	416.0
		10	476.0	509.3
		20	544.0	554.6
		30	612.0	662.6
		Mean		535.6
	CD	0	384.0	396.0
		10	448.0	466.0
		20	512.0	528.0
		30	569.3	588.0
		Mean		494.5
	IF	0	432.0	412.0
		150	504.0	492.6
300		576.0	642.3	
450		642.6	651.6	
Mean			549.6	
BR9922-DMRSF ₂	PM	0	336.0	400.0
		10	392.0	416.0
		20	448.0	484.0
		30	504.0	528.0
	Mean		457.0	
	CD	0	288.0	404.6
		10	336.0	418.6
20		384.0	465.3	

		30	432.0	526.0
		Mean		453.6
	IF	0	360.0	412.0
		150	420.0	492.6
		300	480.0	642.3
		450	540.0	651.6
		Mean		549.6
Agbor Local	PM	0	312.0	356.6
		10	364.0	375.3
		20	416.0	428.0
		30	468.0	476.6
		Mean		409.1
	CD	0	248.0	380.6
		10	280.0	402.6
		20	304.0	439.3
		30	384.0	471.3
		Mean		423.5
	IF	0	260.0	376.0
		150	303.0	474.6
		300	364.0	480.0
		450	116.0	500.0
		Mean		457.7
Variety			*	*
Manure type			*	*
Rate			*	*
Variety x manure type			Ns	*
Variety x rate			*	*
Manure type x rate			Ns	Ns
Variety x manure type x rate			Ns	Ns

Legend: * = significant at 0.05 level of probability, ns = not significant

Discussion

Soil Physico-chemical Properties of the Experimental Site

The sandy loam texture of the experimental site may be attributed to the Parent Material (PM) from which the soil was formed and the climate of the area. The soil might be formed from sandstone and quartz parent materials. These impart sandy texture to the soils. The high sand content of the soil could be attributed to high content of quartz in the material (Brady and Weils, 1999). The weak acid nature of the soil of the area may be traced to the marked leaching of exchangeable bases resulting from the high rainfall associated with the environment and the dissociation of strong and functional group in the organic matter. This is

in harmony with the findings of Omokri *et al.* (2007). The low organic matter status of the experimental site could be attributed to the rapid decomposition of organic matter due to high solar radiation and moisture, this favours optimum microbial activities in the soil. It could also be attributed to the annual seasonal bush burning which tend to deplete organic matter accumulation in the soil (Landor, 1991).

The low level of total nitrogen could be due to high temperature. It could also be attributed to leaching of nitrate by torrential rainfall prevalent in the environment (Olatunji *et al.*, 2007). The high level of Phosphorus may be attributed to either of these reasons: (i) history of land use and cultural practices associated with the land use (that is, cropping of crops that do

not take much P nutrient from the soil and the application of P organic fertilizers (Nnaji *et al.*, 2002, Nnaji, 2008). (ii) The parent material from which the soil was formed may be rich in P minerals (Brady and Weils, 1999). (iii) The soil may not be highly acidic as to cause high level of fixation (Brady and Weils, 1999, Isirimah *et al.*, 2003 and Omokri *et al.*, 2007). The low values of exchangeable cations may be attributed to the leaching of bases from the solum due to high rainfall characteristics of the area. The low action exchange capacity could be attributed to the PM from which the soil was formed, and low organic matter (OM) content of the soil. The PM from which the soil was formed may be poor in basic nutrients. Nnaji *et al.* (2002) noted that soils of the study area were dominated by Fe oxide and Kaolinites. These clay minerals are low in basic cations (Brady and Weils, 1999). The exchangeable acidity was low possibly because of the cultural practices associated with the land use (that is, previous use of amendments to improve soil condition and enhance crop yield. The results, generally, are in harmony with the findings of Olatunji *et al.* (2007) and the results of soil fertility evaluation in the region. It is also consistent with the findings of Nnaji *et al.* (2002), Nnaji, 2008) which reported that the history of land use and cultural practices affect soil conditions and crop productivity.

Nutrient Content (% dry matter) of Organic Manure used in the Study

The values of N, P and K were higher in poultry manure than in cattle dung used in the study possibly because poultry manure, especially those produced in deep litter or battery cage house, have more concentrated nutrient content compared with other types of animal manure. This is similar to the findings of Sharpley and Smith (1995) and Brady and Weils (1999) who reported that among the different sources of organic manure which have been used in crop production, poultry manure was found to be the most concentrated in terms of nutrient content. It is also in harmony with the findings of Subedi and Gerung (1991) and DIPA (2006) who reported that poultry manure has higher levels of N, P and K than cattle dung. It is also consistent with the findings if Ibeawuchi *et al.* (2007) who reported higher levels of N, P and K in plots treated with poultry manure than in plots treated with other nutrient sources.

Effect of Variety, Organic Manure and Inorganic Fertilizer on Number of Grains/cob of Maize

Hybrid variety 9022-13 had higher number of grains/cob than open-pollinated variety BR9022-DMRSRF₂ and Agbor local possibly because hybrid maize varieties possess outstanding quality with respect of high grain yield. This is consistent with the findings of Odeleye and Odeleye (2001) and Sajjan *et al.* (2002) who reported that fruit production, grain yield and other yield components are usually influenced by genetic quality of individual variety. It is also in consonance with the findings of Olakojo *et al.* (1998) who reported that hybrid maize varieties have yield advantage of 25 to 50% over the best open-pollinated variety in Nigeria. Plants that received poultry manure had higher number of grains/cob than plants that received cattle dung probably because poultry manure must have increased the carbon content, water holding capacity, aggregation of soil, and decreased the bulk density of the soil. This is similar to the findings of Egerszegi (1990) who reported that poultry manure increased the carbon content, water holding capacity, aggregation of soil and decreased bulk density of soil where it was applied. Poultry manure could have also increased the water soluble and exchangeable potassium and magnesium, which enhanced yield as reported by Jackson *et al.* (1999). The superiority of plants that received inorganic fertilizer over other plants with respect to higher number of grain/cob could be attributed to better yield enhancement credited to inorganic fertilizers over other nutrient sources. This is similar to the findings of Ayodele (1993) and Giller (2003) who reported that inorganic fertilizers influenced crop quality and improved yield over 50% than their unfertilized counterparts. It is also consistent with the findings and report of Akintoye and Olufolaji (2005) on *Cayene* pepper. Higher rates of inorganic fertilizers or organic manure produced better response with respect to higher number of grains/cob possibly because higher rates satisfied the crop's requirement. This is similar to the findings of Olarewaju and Isma (1990) and Ayodele (1993) who suggested that accurate rate and type of fertilizer should be applied using the best method at the most appropriate time.

Conclusion and Recommendations

This study was carried out to evaluate the effects of variety and fertilizers on number of grains/cob of maize in Asaba area of Delta State. It was a factorial experiment carried out in a Randomized Complete Block Design (RCBD) with three replicates. Four different rates of poultry manure, cattle dung and inorganic fertilizer NPK 20:10:10 were applied to three maize varieties sown at 75 cm x 15 cm plant spacing. The results obtained indicated that hybrid variety 9022-13 was superior in number of grains/cob of maize and that plants which received inorganic fertilizer application rate of 450 kg ha^{-1} had the highest number of grains/cob. Based on the findings of the study, it was recommended that (i) Hybrid maize variety 9022-13 be grown in the study area (ii) farmers who prefer mineral fertilizer should apply 450 kg ha^{-1} to enhance number of grains/cob of maize in Asaba area of Delta State.

References

- Adekunle, I. O., Akirinade, O. E. and Azeaz, J. O. (2005).** Influence of combined application of cattle manure and NPK fertilizer on soil chemical properties, growth and yield of Okra (*Abelmoschus esculentum*) in Alfisol. *Proceedings of the 29th Annual Conference of the Soil Science Society of Nigeria, December 6-10, 2004, University of Agriculture, Abeokuta, Nigeria.* Pp. 143-146.
- Adepetu, J. A. (1997).** Soil and Nigeria food security. Inaugural Lecture series 119, Obafemi Awolowo University, Ile-Ife. Nigeria p.19.
- Agbato, S. O. (2003).** Principles and Practices of crop production. Odumatt press publisher, Oyo, pp. 57-62.
- Akintola, F. E. (1997).** Maize production: constraints and Improvement in Nigeria. The maize Association of Nigeria Ibadan. Pp. 223-232.
- Akintoye, H. A. and Olufolaji, A. O. (2005).** Effect of different levels of organic mineral fertilizer on the growth and yield of Cayenne pepper grown in South Western Nigeria. In HORSTON Abstracts 2005 of 23rd Annual conference held on September 16-22 at Rivers State p. 35.
- Allessi, J. and Power, J. F. (2004).** Effect of plant population, row spacing and relative maturity on dry land corn in the Northern plains. *Agronomy Journal*, 66(2), 316-319.
- Ayodele, O. J. (1993).** Further yield responses of tomatoes (*Lycopersicum esculentum*) to fertilizer application *Research Bulletin* No. 16. national Horticultural Research Institute, Ibadan, Nigeria.
- Azam, S., Ali, M., Amin, M., Bibi, S. and Arif, M. (2007).** Effect of plant population on maize hybrids. *Journal of Agricultural and Biological Science*, 2(1), 104 – 111.
- Bowles, J. E. (1992).** Engineering properties of soils and their management. 4th (ed). McGraw Hill Publishers, Boston, U.S.A. p 241.
- Brady, C. and Weils, R. R. (1999).** Nature and properties of Soil Twelfth Edition, Prentice Hall, New Delhi pp 74 –114.
- DIPA (2006).** Handbook of Agriculture: facts and figures for farmers, students and all interested in farming. Directorate of Information and Publications of Agriculture. Indian Council of Agricultural Research, New Delhi, p. 435.
- Dut, S. (2005).** A Handbook of Agriculture. ABD Publishers, India. Pp 116-118.
- Egerszegi, E. (1990).** Effect of sewage sludge and compost applied to the soil on some physical and chemical properties. *Journal of Environmental Quality*, 15,122-127.
- FAO (2002).** Fertilizer and the future. *IFA/FAO Agriculture Conference on Global food security and the role of Sustainability Fertilization.* Rome, Italy. 16th-20th March, 2003, pp 1-2.
- Fayenisin, O. (1993).** Search for Improved maize varieties for farmers in Nigeria. 3rd National Workshop of Maize Centre. NASPP, Ibadan December, 6th - 10th 1976.
- Giller, K. E. (2003).** Targeting management of organic resources and mineral fertilizer. Can we match scientist fantasies with farmer realities. In integrated plant nutrient management in Sub-Saharan Africa, from concept to practice (B. Vanlauwe J. Diels, N. Sanginga, R. Merelox (eds) pp. 211-217.
- Ibeawuchi, I. I., Opara, F. A, Tom, C. T. and Obiefuna, J. C. (2007).** Graded replacement of inorganic with organic manure for sustainable maize production in Owerri Imo State, Nigeria. *Life Science Journal*, 4(2), 82-87.
- Iken, J. E. and Anusa, A. (2004).** Maize Research and Production in Nigeria. *African Journal of Biotechnology*. 3(6), 302-307.
- Iken, J. E., Anusa, A. and Obalaju, V. O. (2001).** Nutrient Composition and Weight Evaluation of some Newly Developed maize

- Varieties in Nigeria. *Journal of Food Technology, Kenya*, 7, 25-28.
- Isirimah, N. O., Dickson, A. A. and Igwe, C. (2003).** Introductory soil Chemistry and Biology for Agriculture and Biotechnology. Asia International Publishers Limited, Port-Harcourt, Nigeria. Pp 103.
- Ismail, A. S., El-Schaay, A. S., Salehu, S. A., and Abdel-Wahab, A. F. (1990).** Effect of application of mineral and organic amendment of nodulation of cowpea growth and certain chemical properties of calcareous soil. *Ann. Agric. Sci. (Special Ed)*, 23-39.
- Jackson, H. (1999).** Land disposal of broiler litter changes in soil potassium, calcium and magnesium. *Journal of Environment Quality*, 4, 202-206.
- Juo, A. S. R. (1981).** Mineralogy of Acid Sands of Southern Nigeria. Monograph N.O.1. Soil Science Society of Nigeria Pp. 19-26.
- Kim, S. K. (1997).** Achievement, challenges and future direction of hybrid maize research and product in B. Badu-Apraku, Akoroda M.O., Oedraw M. and Quin F.M(eds). *Proceedings of Required Maize Workshop May 99-june 2, 1995. IITA Cotonou, Benin Republic.*
- Landor, J. R. (1991).** Brooker tropical soil manual. A handbook for soil survey in the tropics and subtropics. Longman Group. England. 106-144.
- Mclean, E. O. (1982).** Aluminum. In C.A. Black, (ed) Methods of soil analysis part 2. American Society of Agronomy, ASA Monograph 9, Pp 978-998.
- Meteorological Office Asaba (2003).** Weather Characteristics of Delta State of Nigeria. Unpublished Vol. 2 p 2-21.
- Movahedi, S. A. R, and Cook, H. F. (2000).** Influence of municipal compost on temperature, water, nutrient status and the yield of maize, In a Temperate Soil. *Soil use and Management*, 16(3), 215-221.
- Nest-Nigerian Environment Study Action Team (1991).** Nigerian threatened environment: A National Profile, p 288.
- Nnaji, G. U. (2008).** Fertility status of some soils in Isoko South Local Government Area of Delta State. *Proceedings of the Annual Conference of the Agricultural Society of Nigeria* held n Ebonyi State University, Abakiliki Pp 515-519.
- Nnaji, G. U., Asadu and Mbagwu, J. S. C. (2002).** Evaluation of the physico-Chemical properties of soils under selected agricultural land utilization types. *Journal of Tropical Agriculture, Food, Environment and Extension* 3, 27-33.
- Odeleye F. O. and Odeyeye, M. O. (2001).** Evaluation of morphological and agronomic characteristics of two exolic and two adapted varieties of tomato (*Lycopersicon esculentum*) in South West Nigeria. *Proceedings of the 19th Annual Conference of HORTSON*. 1,140-145.
- Olakojo, S. A., Ogundodede, B. A. and Kogbe J. O. S (1998).** Evaluation of Maize (*Zea mays*) Top crosses in a Rainforest Location. *Biose res. Comm.* 11(2), 141-146.
- Olanikan, P. C. (2006).** Organic manures as Soil Amendment in Eroded Tropical Soil of South Western Nigeria. *Soil and Nutrition. Journal of Tropical Soils*, 5, 11-18.
- Olarewaju, J. U. and Isma, F. (1990).** Irrigated pepper production as influenced by plant population and nitrogen nutrition at Samand, Nigeria. *Paper presented at 26th Annual conference of Agriculture, Markudi, 2-5 September, 1990.*
- Olatunji, U. S. A., Ayuba, A. and Oboh, V. U. (2007).** Growth and yield of Okra and Tomato as Affected by pig dung and other organic manures. *Proceedings of the 30th Annual Conference of the soil science society of Nigeria 5-9th December, 2005, Makurdi Nigeria.*
- Olayinka, A. (1996).** Carbon Mineralization from Poultry Manure Straw Sawdust amended alfisol. *Ife Journal of Agriculture* 1 and 2. (18,) 26-36.
- Olayinka, A., Adentunyi, A. and Adebayo, A. (1998).** Effect of organic amendment on nodulation and nitrogen fixation of cowpea. *Journal of Plant Nutrition*, 21(11), 2455-2464.
- Omokri, J. I., Nnaji, G. U. and Uyovbisere, E. O. (2007).** Assessment of the physical properties of Orogo basin and the surrounding soils in in Agbor, Delta State. *Proceedings of the 31st Conference of Soil Science Society of Nigeria* held on the 13th – 17th November, 2006 at Ahmmadu Bello University, Zaria, Nigeria Pp 358-362.
- Reijntjes, C., Hoverkork, B. and Water-Bayer, A (1992).** An introduction to Low External-Input and Sustainable Agriculture. John Wiley and Sons. London. p. 340.
- Sajjan A. S., Shekhargounda, M. and Badanur (2002).** Influence of data of sowing, spacing and levels of nitrogen on yield attributes and seed yield of Okro. *Ikamataka Journal of Agricultural Science*, 15(2), 267-274.

- Sharpley, A. N and Smith, S. J. (1995).** Nitrogen and phosphorus forms in soil receiving manure. *Soil Science*, 159, 253-258.
- Sobulo, R. A. and Babalola (1992).** Improved organic fertilizer and soil condition. *In Toward Efficiency Fertilizer use in Nigeria. Federal Ministry of Agriculture, Water Resources and Rural Development.* Lagos pp. 90-110.
- Sonetra, S., Borin, K. and Preston, T. R. (2002).** Waste water from rubber processing as fertilizer for water spinach and forage cassava. www.utafoundation.org/utacanbod/msc99thes/so_nintro.htm.
- Stefan, T. (2003).** Organic Agriculture: Sustainability market policies. CAB Organization for Economic Co-operation and Development p. 95.
- Subedi, K. D. and Gurung, G. (1991).** Soil fertilizer thrust towards sustainable agriculture: experience of Lumile Regional Agricultural Research centre. *In: Soil Fertility and Erosion Issues in the Middle Mountains of Nepal* pp. 81-82 IDRC, Ottawa, Canada.
- Wahua, T. A. T (1999).** Applied statistics for scientific studies. Afrika-Link Books, Nigeria, pp. 250 -287.
- Walkely, A. and Black, I. A. (1945).** An Examination of Detrigareff methods for determining soil organic matter and proposed modification of the chronic and titration methods. *Soil Science*, 37, 29-38.
- Zeidan, M. S., Amany, A. and Balor El-Kramany, M. F. (2006).** Effect of N-Fertilizer and lant Density on Yield and Quality of maize in Sandy Soil. *Research Journal of Agriculture and Biological Sciences*, 2(4), 156-161.